

Submitted to Ap. J.

OBSERVATIONAL EVIDENCE OF THE FORMATION OF CYANOPOLYYNES IN CRL618 THROUGH THE POLIMERIZATION OF HCN

Juan R. Pardo, José Cernicharo & Javier R. Goicoechea¹

*IEM - Departamento de Astrofísica Molecular e Infrarroja, CSIC, Serrano 121, E-28006
Madrid, Spain.*

¹*Present address: LERMA-École Normale Supérieure (UMR 8112 du CNRS), 24 Rue
Lhomond, 75231 Paris, France.*

pardo,cerni,javier@damir.iem.csic.es

ABSTRACT

The abundance ratio of consecutive members of the cyanopolyynes family has been explored in CRL618 using data acquired in a complete line survey covering the frequency range 81-356 GHz. The J_{up} range explored for the different molecules is the following: 1 to 4 for HCN and HNC, 9 to 39 for HC₃N, 31 to 133 for HC₅N, and 72 to 85 for HC₇N (not detected beyond $J_{up}=85$). The lowest vibrationally excited state of HC₇N (ν_{15} at 62 cm⁻¹) has been tentatively detected. Data analysis has been performed by extending our previous geometrical and radiative transfer model of the slowly expanding envelope (SEE) surrounding the compact central continuum source of CRL 618, that was established from the study of rotational lines in several vibrationally excited states of HC₃N. The new lines analyzed here require to model the high velocity wind (HVW) component and the colder circumstellar gas, remnant of the AGB phase of CRL618. The derived HC₃N/HC₅N and HC₅N/HC₇N abundance ratios from this set of uniformly calibrated lines are between ~ 3 and 6 in the different regions, similar to standard values in the CSM and ISM, and consistent with previous estimates obtained from ISO observations and chemical models. However, the abundance ratios of HC₃N, HC₅N and HC₇N with respect to HCN are at least two orders of magnitude larger than those typical for AGB C-rich stars, such as IRC+10216. This fact indicates that, in the short transition toward the Planetary Nebula phase, HCN is quickly reprocessed into longer cyanopolyne chains. A similar behavior was previously found in this object for the polyacetylenic chains (C_{2n}H₂).

Subject headings: stars: post-AGB-stars: carbon-rich - stars: circumstellar matter - stars: individual: CRL618 - ISM: molecules - radio lines: stars

1. Introduction

CRL 618 is a C-rich protoplanetary nebula (PPNe). Its chemical richness and its complex morphology with a central B0 star surrounded by an ultracompact HII region, and a thick molecular and dusty envelope (Bujarrabal et al. 1988) with different outflows (Cernicharo et al. 1989, hereafter CER89), are now well known thanks to many detailed observational studies. Detailed chemical models for this object have been developed (Cernicharo 2004, hereafter CER04) indicating the rapid evolution of the central star and its influence on the circumstellar ejected material. Changes seem to occur at short -almost human- time scales. The most recent observational results on the morphology are the interferometric observations presented in Sánchez-Contreras & Sahai, 2004. Concerning the chemical composition, a line survey from 80 to 275 GHz has been recently completed (Cernicharo et al. 2005).

After the study of the physical conditions in the slowly expanding envelope (SEE) around the ultracompact HII region of CRL 618 using vibrationally excited states of HC_3N (Pardo et al. 2004, hereafter PAR04), we focus now on the entire cyanopolyynes family: HCN, HNC, HC_3N , HC_5N and HC_7N (longer members have not been detected in our survey).

HC_5N is the second molecule, after HC_3N , in terms of total number of lines detected in the millimeter spectrum of CRL 618. It is a linear molecule with a rotational constant of 1331.330 MHz and a dipole moment of 4.33 Debyes, first discovered in space by Little et al. (1978). The two lowest energy doubly degenerated bending modes (ν_{11} and ν_{10}) have an energy of 105 and 230 cm^{-1} respectively (Yamada et al., 2004). On the other hand, HC_7N was first discovered in space by Kroto et al. (1978). Its rotational constant and dipole moment are 564.0011 MHz and 4.82 Debyes respectively. $\text{HC}_3\text{N}/\text{HC}_5\text{N}$ and $\text{HC}_5\text{N}/\text{HC}_7\text{N}$ abundance ratios between 1.4 to 3.0 were found in molecular clouds since the earliest studies (Snell et al. 1981). Finally, HCN, with a dipole moment and rotational constant of 2.98 Debyes and 44315.976 MHz respectively, is one of the most abundant molecules in the interstellar and circumstellar media. In many cases HCN lines are optically thick so that observations of isotopologues or vibrationally excited states are preferred to probe molecular clouds and circumstellar envelopes. **The other two molecules discussed in this paper have the following dipole moments and rotational constants: 3.73 Debyes and 4549.059 MHz (HC_3N), and 2.98 Debyes and 45331.7845 MHz (HNC).**

The goal of the study conducted in this paper is to get abundance ratios of the different members of the cyanopolyynes family in CRL 618 and to compare them with those observed in prototypical AGB objects such as IRC+10216. In addition, by using new polar species we aim at improving and extending our previous model of CRL618 (see PAR04). This is another step toward a final model for the whole millimeter wave spectrum of this source. The observational procedure and SEE model developments were described in detail in PAR04, we only give here a short summary in sections 2 and 3. The analysis of the HCN, **HNC**, $v=0$ HC₃N, HC₅N, and HC₇N data, with the extension of the model, is presented in section 4. Comparisons of the results with chemical model predictions, and other discussions, are presented in section 5. The work is summarized in section 6.

2. Observations

The observations presented in this paper are part of two line surveys of CRL618. The first one, now complete, has been carried out with the IRAM-30m telescope from 1994 to 2002 (Cernicharo et al. 2005, hereafter CER05) with a frequency coverage of 81-279 GHz except for frequencies with high atmospheric opacity around 119 and 183 GHz. The second one, still on-going but complete for the species we focus on in this work, is performed with the CSO telescope between 280 and 356 GHz. **For both instruments the frequency resolution used has been around 1 MHz with ~ 0.5 GHz wide spectrometers.** The J_{up} range covered for rotational transitions of HCN **and HNC** is 1 to 4, for HC₃N 9 to 39, for HC₅N 31 to 133, and for HC₇N it is 71 to 316, although the lines are not detected beyond $J_{up}=85$. See PAR04 for details on the observational procedures at the two facilities.

3. Previous Model Summary

A model has been developed to study the physical conditions that explain the whole observed millimeter wave spectrum of CRL618. The precise knowledge of the spectral behavior of the continuum emission (see PAR04) is paramount since it plays a very important role in the observed line profiles (specially in the emission-to-absorption line ratio). The model consists of a gas envelope around a central continuum source, considered spherical with size and effective temperature adjusted to fit the continuum IRAM-30m data (PAR04) and assumed to keep the same spectral index in the range 280-356 GHz (CSO observations). The description of the gas component allows different shells with different physical conditions to be included in the model. The geometry is not restricted to be spherical. Elongated shells with both radial and azimuthal velocity components can be defined.

Table 1: Observational parameters (velocity centroid, width and integrated area) obtained from gaussian fits of the absorption and emission components of selected HC₅N lines in **CRL618**. See **beginning of** section 4 for **more details**.

J_{up}	v state v	ν (GHz)	Absorption			Emission		
			v (km·s ⁻¹)	Δv (km·s ⁻¹)	A K·km·s ⁻¹	v (km·s ⁻¹)	Δv (km·s ⁻¹)	A (K·km·s ⁻¹)
33	$v=0$	87.86363	-20.3	18.9	2.7	-43.3	23.3	-0.87
33	$\nu_{11} 1^-$	88.00874	-20.4	10.0	0.29	-33.1	8.6	-0.29
33	$\nu_{11} 1^+$	88.08534	-22.6	23.2	0.54	-34.7	14.1	-0.55
33	$2\nu_{11} 0$	88.22163	-21.1	6.8	0.09	-30.5	16.9	-0.17
33	$2\nu_{11} 2^-$	88.23055	-24.8	-	-	-31.2	-	-
33	$2\nu_{11} 2^+$	88.24034	-23.2	5.1	0.09	-33.5	5.9	-0.08
33	$\nu_{10} 1^+$	88.04187				-34.7	6.7	-0.05
33	$3\nu_{11} 1^-$	88.33509				-31.2	4.9	-0.05
33	$3\nu_{11} 3^+$	88.41807				-28.5	7.2	-0.16
33	$3\nu_{11} 1^+$	88.48921				-31.4	7.2	-0.12
34	$v=0$	90.52589	-21.1	19.1	4.0	-38.4	34.1	-1.9
35	$v=0$	93.18813	-21.1	19.3	4.2	-37.3	32.7	-2.6
35	$\nu_{11} 1^-$	93.34201	-22.5	8.4	0.49	-33.0	18.7	-0.89
35	$\nu_{11} 1^+$	93.42323	-23.2	22.2	0.88	-33.5	12.9	-0.69
35	$2\nu_{11} 0$	93.56650	-21.7			-32.0	8.2	-0.30
52	$v=0$	138.44167	-21.3	19.8	3.6	-38.0	17.7	-1.3
58	$v=0$	154.41110	-21.3	14.8	2.8	-38.0	20.8	-1.2
59	$v=0$	157.07253	-21.3	16.1	2.9	-38.0	19.9	-1.2
59	$\nu_{11} 1^-$	157.33140	-23.8	13.3	1.6	-28.3	17.6	-1.2
59	$\nu_{11} 1^+$	157.46775	-23.7	14.5	0.92	-31.8	8.6	-0.47
59	$2\nu_{11} 0$	157.67636	-19.1	16.6	0.43	-29.9	6.8	-0.19
59	$2\nu_{11} 2^-$	157.72678	-21.0	14.0	0.47	-31.2	9.7	-0.28
59	$\nu_{10} 1^+$	157.39079	-20.0	13.2	0.32			
96	$v=0$	255.50940	-24.6	10.1	0.94			
96	$\nu_{11} 1^-$	255.92839	-23.6	6.1	0.27			
98	$v=0$	260.82794	-21.6	10.9	0.88			

Since the observations reveal that different gas regions are traced by different molecular species, we proceed in our analysis in different steps. The first one **has been to** study those species arising from the inner $\sim 1.5''$ SEE, with outflow velocities of ~ 10 to $15 \text{ km}\cdot\text{s}^{-1}$, turbulence velocities around $3.5 \text{ km}\cdot\text{s}^{-1}$, and temperatures in the range 250-275 K. **The full description of the SEE model is given in PAR04. The extension of the SEE model to include both the colder and outer circumstellar gas (necessary to explain the rotational lines in the ground vibrational state of HC_3N , HC_5N and HC_7N) and the high velocity wind component (seen in lines of some abundant species, including $v=0 \text{ HC}_3\text{N}$) is one of the targets of this work. This is discussed in detail in the next section.**

4. Analysis and Discussion

The first step of the analysis has been to perform gaussian fits of selected HC_5N lines in the 3, 2, and 1.3 mm windows. A summary **for a set of lines representative of the different vibrational states and frequency ranges** is presented in Table 1 where the areas are expressed in $\text{K}\cdot\text{km}\cdot\text{s}^{-1}$ ($T_A^* \cdot v_{LSR}$), parameters that could not be determined are marked with a dash, and parameters that had to be fixed in order to obtain a reasonable fit are highlighted in bold face. Also, when the velocity, width and area of one component are left blank in the table it means that the component is not detected in the particular transition. These observational results already reveal an interesting behavior that is crucial for the analysis: lines in the $v=0$ state have absorption and emission components that are in general 15 to $25 \text{ km}\cdot\text{s}^{-1}$ wide at 3 and 2 mm, with the absorption centered at around -37 to $-42 \text{ km}\cdot\text{s}^{-1}$, whereas lines in vibrationally excited states, in the same spectral region, appear narrower (~ 7 to $15 \text{ km}\cdot\text{s}^{-1}$) with the absorption component (sometimes the only one detected) centered at ~ -28 to $-33 \text{ km}\cdot\text{s}^{-1}$. These differences tend to disappear as frequency increases and the line profiles in the $v=0$ state are dominated by the emission component only. These facts suggest that the rotational lines in the ground state of HC_5N are mostly formed in a region with quite different physical conditions than the one responsible for the lines in vibrationally excited states. This is carefully explored in this section.

4.1. Vibrationally excited HC_5N

The latest laboratory data on the ν_{11} and ν_{10} vibrational states of HC_5N , their overtones and their combinations, have been presented by Yamada et al. (2004). In our line survey, we have detected rotational lines in the following vibrationally excited states of HC_5N (most

of them for the first time in space): ν_{11} , $2\nu_{11}$, ν_{10} , and $3\nu_{11}$, at energies of 105, 210, 230, and 315 cm^{-1} respectively. Since these lines also should arise from the SEE analyzed in PAR04, we have just tried to find the $\text{HC}_3\text{N}/\text{HC}_5\text{N}$ abundance ratio that best fits the observations, keeping the same SEE physical parameters of PAR04. The result is $\text{HC}_3\text{N}/\text{HC}_5\text{N} \sim 3 \pm 1$ (see Fig. 1). This result confirms previous estimates made from vibrational bands of both species seen in absorption by ISO around 15 μm (Cernicharo et al. 2001). Obviously, the fit cannot be perfect for so many lines. In particular, Fig. 1 shows that some absorption exists between -50 and -40 $\text{km}\cdot\text{s}^{-1}$ for $J_{up}=35$ in the ν_{11} state that is not reproduced by the model. This can be due to a fraction of a higher velocity outflow that intersects the line of sight toward the continuum source that has not been considered in the model. This absorption component is otherwise clearly visible in the rotational lines inside the ν_7 state of HC_3N , but does not affect the other lines shown in Fig. 1. The detections in the HC_5N vibrational states extend from $J_{up}=31$ to $J_{up}=104$ for ν_{11} , $J_{up}\sim 87$ for $2\nu_{11}$, $J_{up}\sim 77$ for ν_{10} , and $J_{up}\sim 53$ for $3\nu_{11}$. The fact that for HC_3N we detect rotational lines from vibrational states a factor 3-4 higher in energy than for HC_5N is consistent, for the sensitivity of the survey, with the derived $\text{HC}_3\text{N}/\text{HC}_5\text{N}$ abundance ratio, and the dipole moment and partition function ratios of both species.

4.2. $^{12}\text{C}/^{13}\text{C}$ ratio in the SEE

The $^{12}\text{C}/^{13}\text{C}$ isotopic ratio in the SEE can be derived using HCC^{13}CN , HC^{13}CCN and H^{13}CCCN in the ν_7 state. The ν_6 state is also detected but the signal-to-noise ratio is too low for any estimation. Figure 2 shows the results of the SEE model for $(^{12}\text{C}/^{13}\text{C})=15$ (PAR04). We estimate an error bar for this value of $\sim \pm 2$. Kahane et al. (1992) gave lower limits for this ratio of 18 and 3.2 depending on which transition ($J=1-0$ or $J=2-1$) of both ^{12}CO and ^{13}CO was used. Our estimate gives a much better constraint, and restricted to the SEE region, because it relies upon tens of lines without the saturation problems of CO.

4.3. Ground state HC_5N

Using the result found in 4.1, we have run the model for the detected rotational lines of HC_5N in its ground vibrational state. From these calculations it becomes clear that the SEE alone cannot account for these lines. A significant contribution from gas at lower temperatures must exist. **In the individual lines, there is no significant evidence of the high velocity wind (HVW) seen in the most abundant molecules (see below).** Therefore, we have expanded the model presented in PAR04 to include a cold circumstellar shell (CCS),

created during the AGB phase (see figure 3). We have not simply considered a spherical geometry for this cold component in order to account for the optical/infrared images of the CSE that show that the envelope has a bipolar structure. This was also pointed out for the molecular gas in CER89 and Neri et al. (1992). The parameters that define the CCS in our model are:

- R_{CCS1} and R_{CCS2} : Inner and outer boundaries of the shell (angles in arcseconds).
- α_T : Truncation angle (degrees), used to take into account the fact that this shell needs to be divided into at least two zones because the molecular density is probably larger in the polar regions along the axis of the high velocity wind. This fact is revealed by the ratio between emission and absorption in the profiles of the analyzed $v=0$ HC₅N rotational lines.
- $T_{rot,X}^{CCS}$: Rotational temperature (K) for species X in the CCS.
- [HC₃N] (at $R=R_{CCS1}$): Number density of HC₃N at the inner part of the shell for $\alpha > \alpha_T$ (for $\alpha < \alpha_T$ the density will be multiplied by a factor ρ_{ccs}). HC₃N will always be our reference molecule for the models of CRL 618. In each region we will consider a constant ratio with respect to it for each molecule.
- d_{CCS} : The radial distribution of HC₃N abundance is considered to vary as $[HC_3N_R] = [HC_3N_{R_{CCS1}}](R/R_{CCS1})^{d_{CCS}}$ in the CCS.
- v_{CCS} and $v_{turb,CCS}$: radial and turbulence velocity of the gas in the CCS.

Just as we did in PAR04 for the physical parameters of the SEE, we had to run a considerable number of models to find a good fit of the ground state HC₅N lines. Again, the key parameters are the number density, the temperature and the position of the CCS. The position has been fixed between **diameters** 3.0'' and 4.5'' after trying several other positions in our models. Then, the temperature and density profiles can be established by running a grid of models. All the other parameters are then explored to fine tune the shape of the line as J_{up} varies. For example, one of the most remarkable differences in the line profiles from ground to vibrationally excited HC₅N is the position of the maximum depth of the absorption, its width and the absorption/emission ratio. As shown in Figure 1 and table 1, the maximum depth of the absorption for vibrationally excited HC₅N happens at ~ -30 km·s⁻¹ in the 2 and 3 mm windows (as for the lines in vibrationally excited states of HC₃N presented in PAR04), whereas for the lines in the vibrational ground state it happens at around -40 km·s⁻¹ and is generally wider and less deep. The observational differences

have allowed us to determine the kinematic conditions in the CCS. The parameters of the best simulation (Fig. 4) are shown on Table 2. The rotational temperature is in the range 50-70 K. In fact, a simulation with $T_{rot}=50$ K provides a better fit for lines below $J_{up} \sim 55$, whereas $T_{rot}=70$ K provides a better fit for the range $J_{up} \sim 55-80$. The simulation shown in Fig. 4 is a compromise between the two cases ($T_{rot}=60$ K). Above $J_{up} \sim 80$, the line profiles are almost exclusively determined by the SEE region. The error bar for the $[HC_3N]/[HC_5N] = 3$ ratio in the CCS can again be estimated to be ± 1 . The same absorption at $v_{LSR} < -40$ km.s⁻¹ reported in section 4.1, and not perfectly modeled here, is seen in some $v=0$ HC₅N spectra. Absorptions centered at ~ -40 km.s⁻¹ have previously been reported in this object in molecules such as CO (Herpin et al. 2002), HCN (CER89 and Neri et al. 1992), HCO⁺ (CER89 and Sánchez-Contreras & Sahai, 2004), HC₃N (CER89, PAR04), and CS (Hajian et al. 1995). There is general agreement to assign that absorption to the AGB circumstellar envelope.

Sánchez-Contreras & Sahai (2004) suggest that an elongated structure of $\sim 11''$ diameter with expansion velocity of ~ 17.5 km.s⁻¹ is responsible for part of the emission of the low velocity molecular gas in their HCN J=1-0 and HCO⁺ J=1-0 interferometric maps. This structure is most probably **an extension (due a larger abundance of this molecules in cold regions)** of the CCS we are modeling here for single antenna data. The difference between this larger shell and our CCS is probably due to the different excitation conditions of the low-J lines of HCN and HCO⁺ and the high-J lines of HC₅N explored in this work.

On the basis of the extended model, that now includes two gas regions, we can check for the presence of longer cyanopolynes rather easily because only their abundance ratio with respect to HC₅N or HC₃N is needed to make predictions of their lines and then check against the data (assuming the same rotational temperature). This is presented in the following subsections.

4.4. Isotopic HC₅N and ¹²C/¹³C ratio in the CCS

We have looked at the possible detection of the ¹³C substituted species of HC₅N by comparing the observations at the frequencies of their rotational lines in the ground vibrational state with the extended model results considering a ¹²C/¹³C isotopic ratio of 15 in the SEE (as derived in section 4.2) and also in the CCS. The simulations indicate that this value is an upper limit for the CCS. This is important to later evaluate the HCN and HNC abundances from H¹³CN and ¹³C (section 4.6).

Table 2: Model parameters of the Circumstellar Shell (see figure 3) that provide the best fit to the HC_5N rotational lines in the ground vibrational state (see figure 4).

parameter	value
$2 \cdot R_{CCS1}$ (arcsec)	3.0
$2 \cdot R_{CCS2}$ (arcsec)	4.5
θ_d (deg)	40
α_T (deg)	30
$T_{rot, \text{HC}_{5,7}\text{N}}^{CCS}$ (K)	50-70
$[\text{HC}_3\text{N}]$ (at $R=R_{CCS1}$) (cm^{-3})	0.8
d_{CCS} (no units)	-1.0
ρ_{CCS} (no units)	1/4
$[\text{HC}_3\text{N}]/[\text{HC}_5\text{N}]$	3
$[\text{HC}_5\text{N}]/[\text{HC}_7\text{N}]$	6
v_{CCS} (at $R=R_{CCS1}$) ($\text{km} \cdot \text{s}^{-1}$)	22
$v_{turb, CCS}$ (at $R=R_{CCS1}$) ($\text{km} \cdot \text{s}^{-1}$)	12
v_{CCS} (at $R=R_{CCS2}$) ($\text{km} \cdot \text{s}^{-1}$)	8
$v_{turb, CCS}$ (at $R=R_{CCS2}$) ($\text{km} \cdot \text{s}^{-1}$)	5

4.5. HC₃N in the CCS

The HC₃N lines in vibrational states below $2\nu_7$ show an emission that exceeds what is predicted by the SEE model. Including the CCS component as in section 4.3 provides a first approximation to the detected signal (see figure 5) that is in fact quite good for J_{up} below ~ 20 . However, the HVW component is also visible in the high- J $v=0$ lines and therefore should be included in the model. The high velocity outflow is modeled by just considering a gas component with a given excitation temperature (350 K), average column density ($2.5 \cdot 10^{17}$ cm⁻²), filling factor at a reference frequency ($1.3 \cdot 10^{-3}$ at 90 GHz) and halfwidth velocity outflow (150 km·s⁻¹). The rather high temperature for this component is confirmed by the dominant role that the broad HVW component has at high J 's (see CSO lines in figure 5). More precise details about the structure of the fast outflow are difficult to obtain with the present data. A better constraint on the temperature should be obtained by observing a few $v=0$ HC₃N lines above 400 GHz (on going).

4.6. HCN and HNC emission

At this point of the analysis we have a model that considers the three main regions of the gas emission in the millimeter wave spectrum of CRL618: SEE, CCS, and HVW. We can try to fit the observed lines from other species by adjusting only their ratio to HC₃N in the assumption that the rotational temperature is the same. This approximation will be quite good for those species arising mainly from the SEE, due to the high density of the region (populations are thermalized). It could also be useful for the HVW. However, the low densities imply that the full analysis for all molecular species will be more complex in the CCS.

In this work we will focus only on HCN and HNC. For opacity reasons, we cannot probe the inner SEE and HVW regions using the main isotopologues of both species, specially as J increases. Instead we can use H¹³CN and HN¹³C assuming a SEE ¹²C/¹³C value of 15 (see above) **but even for these isotopologues some contribution from the CCS can be expected. As a result, our simulations considering only the SEE+HVW (see figure 6) for $J_{up} > 2$ provide upper limits for the H¹³CN/HC₃N and HN¹³C/HC₃N ratios in the SEE. The values obtained are about 1/2 and 2/15 respectively which translate into HCN/HC₃N ~ 7.5 and HCN/HC₃N ~ 2.0 .**

To improve the above estimates in the SEE it is better to use vibrationally excited states of HCN and HNC since the CCS contribution should be negligible for them. The lowest one ($v_2=1$), at 712 and 477 cm⁻¹ respectively for HCN and HNC,

is detected in both molecules (l doublets for all rotational transitions except the forbidden $J=1-0$). We have selected those lines that have been observed and are not badly blended with other species to get a better constraint for the above ratios. This analysis (Figure 7) results on $\text{HCN}/\text{HC}_3\text{N} \sim 2$ **and** $\text{HNC}/\text{HC}_3\text{N} \sim 1/3 \text{ to } 1/5$. It is thus demonstrated that much lower ratios of HCN and HNC with respect to longer cyanopolyynes are found in the inner gas of CRL618 (PPNe phase) compared to those found in IRC+10216 (AGB phase). See discussion in section 5.

4.7. Ground state HC_7N

Finally, we have studied the HC_7N rotational lines within the $v=0$ state. This molecule is clearly detected since all transitions between $J_{up}=72$ and $J_{up}=85$ appear above a 3σ level except for three cases due to a coincidence with a much stronger line (see Table 3 and figure 8). Note that most of the HC_7N $v=0$ signal arises from the CCS. We reach this conclusion after running models for which no HC_7N was present in the CCS, and its ratio to HC_5N was set to be $1/3$ in the SEE. In this case, the model predicts some detectable absorption and almost no noticeable emission. The fact that HC_7N is seen almost exclusively in emission with a much larger intensity than in this model, can be easily reproduced if the CCS is included with a $\text{HC}_5\text{N}/\text{HC}_7\text{N}$ abundance ratio around 3 and the same rotational temperature for both species (see figure 8). The model also predicts that above $J_{up}=85$ (at an energy level of 134 cm^{-1}) the signal would be too weak to be detected above the 3σ level of the survey. Similarly, the intensity of lines of HC_9N , with a ratio of $1/3$ or less with respect to HC_7N , is estimated to be well below the detection limit of our observations.

4.8. Vibrationally excited HC_7N

The frequencies of the rotational transitions in the lowest energy bending modes of HC_7N ($\nu_{13}, \nu_{14}, \nu_{15}$) have become available in a recent work by Bizzocchi & Degli Esposti (2004). We have searched for those in the lowest vibrational state $(0,0,1)$ but no individual detection has been obtained. These lines have been modeled by considering computed ab initio values for the $(0,0,1)$, $(0,1,0)$ and $(1,0,0)$ energies: 62 , 163 and 280 cm^{-1} respectively (Botschwina et al. 1997). A tentative detection is obtained when performing the addition in the v_{LSR} scale of the observed signal at the positions of the 1^- and 1^+ $J_{up}=72$ to 79 lines (see figure 8). Five lines in the considered J_{up} range were discarded from the summation because of blending with stronger lines (see Table 3).

Table 3: Detected lines of HC₇N toward CRL 618 (laboratory frequencies in GHz are rounded to 0.1 MHz). **Frequencies** in parenthesis correspond to **co-added lines** in the velocity scale for a tentative detection of ν_{15} (see text).

J_{up}	Vibrational state of HC ₇ N		
	$v=0$	$\nu_{15} 1^-$	$\nu_{15} 1^+$
72	81.2101	(81.3115)	(81.3630)
73	82.3379		(82.4929)
74	83.4656		
75	84.5933	(84.6989)	
76	85.7211	(85.8281)	(85.8824)
77		(86.9572)	(87.0122)
78	87.9765		(88.1421)
79	89.1042	(89.2154)	(89.2719)
80			
81	91.3596		
82	92.4873		
83			
84	94.7426		
85	95.8703		
86			
87			
88			

5. Comparison with chemical models

The abundance ratio for consecutive members of the cyanopolyynes family of molecules in IRC+10216 is ~ 3 (Cernicharo et al. 1987). However, the HCN/HC₃N ratio in the same object is larger than 100. In CRL 618 our results indicate also an abundance ratio near 3 for HC₃N/HC₅N and ~ 6 for HC₅N/HC₇N, but for HCN/HC₃N it is around 2-3. The difference in this ratio at different evolutionary stages of C-rich objects tells us about the chemical processing of the gas when the central star evolves toward the white dwarf stage. CER04 has modeled the role of the strong UV field arising from the hot central star on the chemical abundances of the polyacetylenes, cyanopolyynes and carbon clusters. This work shows that HCN is quickly photodissociated allowing the production of CN which reacts with C₂H₂ and C₂H to produce HC₃N and longer cyanopolyynes, thus explaining the low HCN/HC₃N ratio found in CRL 618. Moreover, this behavior is similar to that found by Cernicharo et al. (2001) for C₄H₂ and C₆H₂. The large abundance of HNC with respect to HCN in CRL618 (HCN/HNC ~ 10 instead of > 200 in IRC+10216, Cernicharo et al. 1987) also points in the same direction. It seems that in a short period of time it could be possible to transform an important fraction of HCN into HC₃N and longer cyanide carbon chains.

6. Summary and future work

We have extended our previous analysis of the gas shells surrounding the protoplanetary nebula CRL618, by studying the pure rotational lines of HC₅N in its fundamental and the lowest 4 vibrationally excited states (first astronomical source in which vibrationally excited HC₅N has been detected), and HC₇N rotational lines in its fundamental vibrational state. We have found that the HC₃N/HC₅N ratio in the innermost slowly expanding envelope is ~ 3 . With this ratio, and all the physical parameters previously derived in PAR04, a good match to the observed HC₅N lines in vibrationally excited states is found. However, the predicted lines for the ground vibrational state are far too weak, suggesting that a more extended and colder gas shell has to be considered. The physical parameters of this envelope, called cold circumstellar shell (CCS) in this work, have been determined. The position of the shell is $\sim 3.0''$ - $4.5''$ from the central star and its excitation temperature is only ~ 60 K (compared to 250-275 K in the innermost SEE). The column density of the CCS in front of the continuum source is negligible compared to that of the SEE. With the CCS derived parameters it is found that the observed HC₇N lines in the ground vibrational state can be well reproduced with a HC₅N/HC₇N abundance ratio of ~ 3 . The isotopic ratio $^{12}\text{C}/^{13}\text{C}$ is found to be ~ 15 in the SEE from the lines of vibrationally excited HC₃N. The same value must be an upper limit in the CCS according to the non detection of ^{13}C substituted isotopologues of HC₅N.

The analysis of the $v=0$ HC_3N lines requires to introduce a third component, the high velocity outflow, although its structure cannot be precisely known. Finally, the ratios $\text{HCN}/\text{HC}_3\text{N}$ and $\text{HNC}/\text{HC}_3\text{N}$ have been derived using the first four rotational transitions of H^{13}CN and HN^{13}C , and some lines of vibrationally excited HCN . The result of a much lower $\text{HCN}/\text{HC}_3\text{N}$ value than in IRC+10216 indicates a processing of HCN into longer cyanopolynes in the protoplanetary nebula stage.

A first examination of lines from all other molecules detected in the survey shows that most of them can be well fitted using the model of the CCS, HVW and SEE gas regions described in this paper and in PAR04, so that a good estimate of their abundances in these regions can be achieved. Therefore, the next step in our analysis of the CRL618 line survey will be to provide a picture of the chemical content in its three main gas components.

We thank the support of the IRAM-30m staff during the long completion of the line survey. CSO operations are supported by U.S. NSF grant AST 22-09008. This work has also been supported by Spanish DGES and PNIE grants ESP2002-01627, AYA2002-10113-E and AYA2003-02785-E.

REFERENCES

- Bizzocchi, L., and Degli Esposti, C, 2004, Ap.J. 614, 518.
- Botschwina, P., Horn, M., Markey, K., & Oswald, R., 1997, Mol. Phys., 92, 381
- Bujarrabal, V., Gomez-Gonzalez, J., Bachiller, R., & Martin-Pintado, J., 1988, A&A, 204, 242.
- Cernicharo, J. Guélin, M., Martin-Pintado, J., Peñalver, J., & Mauersberger, M. 1989, A&A, 222, L1.
- Cernicharo, J., Guélin, M., Menten, K. M., & Walmsley, C. M., 1987, A&A, 181, L1.
- Cernicharo, J., Heras, A. M., Tielens, A.G.G.M., Pardo, J. R., Herpin, F., Guélin, M., Waters, L.B.F.M. 2001, Ap.J. 546, L123.
- Cernicharo, J., 2004, Ap.J., 608, L41.
- Cernicharo, J., & Pardo, J. R., 2005, A&A, In preparation.
- Hajian, A.R., Phillips, J.A., Terzian, Y., 1995, ApJ, 446, 244.

- Herpin, F., Goicoechea, J.R., Pardo, J.R., Cernicharo, J. 2002, Ap.J., 577, 961
- Kahane, C., Cernicharo, J., Gómez-González, J., Guélin, M., 1992, A&A, 256/1, 235.
- Kroto, H.W., Kirby, C., Walton, D.R.M., Avery, L.W., Broten, N.W., MacLeod, J.M., Oka, T., 1978, ApJ, 219, L133.
- Little, L.T., MacDonald, G.H., Riley, P.W., Matheson, D.N., 1978, MNRAS, 183, 45P.
- Neri, R., Garcia-Burillo, S., Guélin, M., Cernicharo, J., Guilloteau, S., Lucas, R., 1992, A&A, 262, 544.
- Pardo, J.R., Cernicharo, J., Goicoechea, J.R., Phillips, T.G., 2004, Ap.J., 615, 495.
- Sánchez-Contreras, C., & Sahai, R., 2004, Ap.J., 602, 960.
- Snell, R.L., Schloerb, F.P., Young, J.S., Hjalmarson, A., Friberg, P., 1981, ApJ, 244, 45S.
- Yamada, K.M.T., Degli Esposti, C., Botschwina, P., Frster, P., Bizzocchi, L., Thorwirth, S., Winnewisser, G., 2004, A&A, 425, 767.




Fig. 1.— Sample of detected lines in vibrationally excited states of HC₅N toward CRL 618. Model results, **shifted by $0.2 \cdot F_c$ (F_c : Continuum Flux) for all vibrational states except for $3\nu_{11}$ ($0.1 \cdot F_c$ shift)**, are shown at the **velocity** resolution of the data ($\Delta v = c \cdot \frac{\Delta \nu}{\nu}$, with $\Delta \nu \simeq 1$ MHz for all spectra) for a HC₃N/HC₅N value of 3. All other physical conditions are as determined in PAR04 in the SEE (the only region considered). The extra feature that appears in the $2\nu_{11}$ $J_{\text{up}}=35$ spectrum is a blending of the $\beta 51$ recombination line of atomic hydrogen plus the $J=83-82$ line of $v=0$ HC₇N.




Fig. 2.— Model predictions considering a $^{12}\text{C}/^{13}\text{C}$ isotopic ratio of 15 in the slowly expanding envelope of CRL 618 (**shifted by $0.2 \cdot F_c$**) compared to the observations for a sample of lines belonging to single ^{13}C substituted HC_3N in the ν_7 state. **The velocity resolutions of model and data are the same ($\Delta v = c \cdot \frac{\Delta \nu}{\nu}$, with $\Delta \nu \simeq 1$ MHz for all spectra and $\nu \simeq J_{\text{up}} \cdot 9.1$ GHz).** Two weak features, between the HCC^{13}CN and $\text{HC}^{13}\text{CCN } \nu_7 1^-$ lines, clearly seen in the spectra with the best signal-to-noise ratio, correspond to the $\nu_6 1^+$ lines of both species. They have also been included in the model. All other extra features that appear in some spectra belong to vibrationally excited states of HC_3N .




Fig. 3.— Model of CRL 618 including a cold circumstellar shell first introduced in our work to reproduce the observed rotational lines of HC_5N in its ground vibrational state. **Details in section 4.3.**




Fig. 4.— (Left) Observed rotational lines of HC_5N in the ground vibrational state compared with the SEE model assuming a $\text{HC}_3\text{N}/\text{HC}_5\text{N}$ value of 3 (dotted histogram, shifted by $1.0 \cdot F_c$) and with the expanded model (solid histogram, shifted by $1.0 \cdot F_c$), that includes the CCS, presented in this paper (Sect. 4, Fig. 3 and Table 2). Both models are basically coincident for J_{up} above ~ 80 . (Right) Results of the CRL 618 model including the SEE and the CCS regions (solid histogram shifted by $0.2 \cdot F_c$) for a sample of lines of ^{13}C substituted HC_5N , compared to the observations. The simulations consider a $^{12}\text{C}/^{13}\text{C}$ value of 15 in both regions. **Velocity resolution of data and model are the same for each plot, as explained in previous figures.**




Fig. 5.— Observed rotational lines of HC_3N in the ground **vibrational** state compared with the SEE+CCS model assuming a $\text{HC}_3\text{N}/\text{HC}_5\text{N}$ value of 3, plus a HVW component according to section 4.5 (**solid histogram shifted by $1.0 \cdot F_c$**). **The model for the SEE region alone is also plotted for comparison (dotted line, shifted by $-1.0 \cdot F_c$)**. Some other lines appear in the data and are also modeled. Most of them are rotational transitions within different vibrational states of HC_3N , HC_5N or their isotopologues, but there are some belonging to other species the discussion of which we leave for a separate work: H_2CO and C_5H in $J_{\text{up}}=16$, and $\text{c-C}_3\text{H}_2$ in $J_{\text{up}}=35$. All observations have been carried out with the IRAM-30m telescope except $J_{\text{up}}=31,35$ (CSO). **Velocity resolutions of data and model are the same as explained in previous figures.**




Fig. 6.— Observed rotational lines of H^{13}CN and HN^{13}C compared with the SEE+HVW model (**shifted by $2.0 \cdot F_c$**) assuming $\text{H}^{13}\text{CN}/\text{HC}_3\text{N}$ and $\text{HN}^{13}\text{C}/\text{HC}_3\text{N}$ values of 1/2 and 2/15 respectively. All observations have been carried out with the IRAM-30m telescope except $J_{up}=4$ (CSO). The very low excitation temperatures of $J_{up} \geq 1, 2$ levels, compared to those in the SEE and HVW regions, results in the poor agreement between data and model for the lines arising from those levels. Nevertheless, the other lines provide a useful **upper limit (the CCS contribution should be important)** about the HCN and HNC abundances in the SEE and HVW regions. **Velocity resolutions of model and data are the same for the different lines, as explained in previous figures.**




Fig. 7.— Observed rotational lines of HCN and HNC in the $v=2$ vibrational state compared with the SEE model (shifted by $2.0 \cdot F_c$) assuming HCN/ HC_3N and HNC/ HC_3N values of 2 and $1/5$ respectively. All observations have been carried out with the IRAM-30m telescope except $J_{\text{up}}=4$ (CSO). Velocity resolutions of data and model are the same as in previous figures.




Fig. 8.— HC_7N ($v=0$) lines observed toward CRL618 compared with model results including the SEE and CCS regions (solid **histogram shifted by $0.075 \cdot F_c$**) and the CCS region alone (dotted line **also shifted by $0.075 \cdot F_c$**). **The considered value of $\text{HC}_5\text{N}/\text{HC}_7\text{N}$ is 3**, which provides a very good fit. The spectrum in the lower-right corner box is the addition of the ν_{15} HC_7N rotational lines listed in table 3 compared to the model results for both the CCS and the SEE. A 4.0 scaling factor has been introduced to keep the same scale of all other spectra.